Mon. Not. R. Astron. Soc. **000**, 1–4 (2006)　　Printed 28 May 2008　　(MN LaTeX style file v2.2)

arXiv:0805.4041v1 [astro-ph] 27 May 2008

The dynamical stability of W Ursae Majoris-type systems

Lifang Li ⋆, and Fenghui Zhang
National Astronomical Observatories/Yunnan Observatory, Chinese Academy of Sciences, P.O.Box 110,
Kunming, Yunnan Province 650011, P. R. China

Accepted yy mm dd. Received yy, mm, dd; in original form 2006 February 13

ABSTRACT

Theoretical study indicates that a contact binary system would merge into a rapidly rotating single star due to tidal instability when the spin angular momentum of the system is more than a third of its orbital angular momentum. Assuming that W UMa contact binary systems rigorously comply with the Roche geometry and the dynamical stability limit is at a contact degree of about 70%, we obtain that W UMa systems might suffer Darwin's instability when their mass ratios are in a region of about 0.076–0.078 and merge into the fast-rotating stars. This suggests that the W UMa systems with mass ratio $q \leq 0.076$ can not be observed. Meanwhile, we find that the observed W UMa systems with a mass ratio of about 0.077, corresponding to a contact degree of about 86% would suffer tidal instability and merge into the single fast-rotating stars. This suggests that the dynamical stability limit for the observed W UMa systems is higher than the theoretical value, implying that the observed systems have probably suffered the loss of angular momentum due to gravitational wave radiation (GR) or magnetic stellar wind (MSW).

Key words: binaries: close – instabilities – stars:blue stragglers – stars: rotation – stars: evolution

1 INTRODUCTION

Interest in W UMa binaries was revived recently with the discovery of large number of new ones among blue stragglers in open and globular clusters (Kaluzny & Shara 1988; Kaluzny 1990; Mateo et al. 1990; Yan & Mateo 1994; Mateo 1996). It appears likely that at least some of blue stragglers are formed by the merging of contact binaries. The models of W UMa systems (Li et al. 2004, 2005) had shown that W UMa systems evolve into contact binaries with extreme mass ratios and then evolve into single, fast-rotating stars (FK Com stars) or blue stragglers due to Darwin's instability when the spin angular momentum of the system is more than a third of its orbital angular momentum (Hut 1980; Eggleton & Kiseleva-Eggleton 2001).

The spin angular momentum has a significant influence on the evolution of W UMa contact binaries, especially for those with extreme mass ratios. Neglecting the spin angular momentum of the small secondary, Rasio (1995) first derived a cutoff mass ratio for contact binary systems and the cutoff mass ratio depends on the structure and the contact degree. Meanwhile, Rasio & Shaprio (1995) identified the dynamical stability limit at a contact degree of about 70% for W UMa systems. Yang (2005b) derived a cutoff mass ratio of about 0.021 from the observed W UMa systems with mass ratios $q < 0.25$. In this work, using the observational data and the theoretical results reported recently, we investigate the dynamical stability limit of W UMa systems, and find that W UMa systems begin merging when their mass ratios are in a region of about 0.076–0.078. This suggests that W UMa systems with mass ratio $q \leq 0.076$ can not be observed. Meanwhile, we estimate that the observed W UMa systems with a minimum mass ratio of about 0.077, corresponding to a contact degree of about 86%, would suffer Darwin's instability and begin merging. This suggests that the minimum mass ratio for the observed W UMa systems is larger than the theoretical value, implying that the observed W UMa systems have suffered the loss of angular momentum loss owing to GR or MSW.

2 DYNAMICAL STABILITY LIMIT

It is well known that a contact binary would suffer Darwin's instability and then coalesce into a fast-rotating single star (including FK Com-type stars and blue stragglers) when the spin angular momentum of the system is more than a third of its orbital angular momentum. The spin angular momentum of a binary star can be expressed as

$$J_{\rm spin} = (k_1^2 M_1 R_1^2 + k_2^2 M_2 R_2^2)\omega_{\rm s}, \qquad (1)$$

⋆ E-mail: gssephd@public.km.yn.cn or llf@ynao.ac.cn where $M_{1,2}$ and $R_{1,2}$ are the masses and radii of the primary and the secondary in solar units, $k_{1,2}$ the ratios of the gyration radii to the stellar radii for both components, and $\omega_{\rm s}$ the spin angular velocity. The orbital angular momentum of the binary system reads

$$J_{\rm orb} = \frac{M_1 M_2}{M_1 + M_2} A^2 \omega_{\rm o}, \tag{2}$$

where $\omega_{\rm o}$ is the orbital angular velocity and A the orbital radius of the binary. We assume that the W UMa systems are in synchronous rotation (i.e. $\omega_{\rm s} = \omega_{\rm o}$) and the ratios of gyration radii to the stellar radii for both components are equal (i.e. $k_1^2 = k_2^2 = k^2$), the ratio of the spin angular momentum to the orbital angular momentum can be written as

$$\begin{aligned} R &= \frac{J_{\rm spin}}{J_{\rm orb}} = \frac{k^2(M_1R_1^2 + M_2R_2^2)}{A^2 M_1 M_2/(M_1+M_2)} \\ &= k^2 \frac{1+q}{q}\Big(\frac{R_1}{A}\Big)^2 \left(1 + q\Big(\frac{R_2}{R_1}\Big)^2\right), \end{aligned} \tag{3}$$

where q $(= M_2/M_1 < 1)$ is the mass ratio of the binary system. If a contact binary system is a marginal contact one in which two components has just filled the inner Roche lobes, following Eggleton (1983), the relative radii of both components of the contact binary read

$$\frac{R_2}{A} = \frac{0.49q^{2/3}}{0.6q^{2/3} + \ln(1 + q^{1/3})}, \tag{4}$$

$$\frac{R_1}{A} = \frac{0.49q^{-2/3}}{0.6q^{-2/3} + \ln(1 + q^{-1/3})}. \tag{5}$$

If the two components of a contact binary system have filled the outer Roche lobes, the relative radii of the components can be expressed by the following equations (Yakut & Eggleton 2005)

$$\frac{R_2}{A} = \frac{0.49q^{2/3} + 0.27q - 0.12q^{4/3}}{0.6q^{2/3} + \ln(1 + q^{1/3})}, \tag{6}$$

$$\frac{R_1}{A} = \frac{0.49q^{-2/3} + 0.15}{0.6q^{-2/3} + \ln(1 + q^{-1/3})}. \tag{7}$$

Using Eqs. (3), (4) and (5) and taking k^2 to be the values of main sequence stars (from 0.05 for no convection stars to 0.21 for fully convective stars), we can obtain a relationship between the angular momentum ratio $R_{\rm i}$ and the mass ratio q for marginal contact systems

$$\begin{aligned} R_{\rm i} = k^2 \frac{1+q}{q}\Big(\frac{0.49q^{-2/3}}{0.6q^{-2/3} + \ln(1+q^{-1/3})}\Big)^2 \times \\ \left(1 + q\Big(\frac{0.6q^{2/3} + q^{4/3}\ln(1+q^{-1/3})}{0.6q^{2/3} + \ln(1+q^{1/3})}\Big)^2\right), \end{aligned} \tag{8}$$

setting equation $R_{\rm i} = 1/3$, and solving (numerically) for q, we obtain the minimum mass ratio $q_{\rm min,in}$ for stability of marginal contact binaries with the different values of k^2 of MS stars and the relation between the minimum mass ratio $q_{\rm min,in}$ and k^2 is shown in Figure 1. Using Eqs. (3), (6) and (7), we obtain a relationship between the angular momentum ratio $R_{\rm o}$ and the mass ratio q for contact systems in which both components have filled the outer Roche lobes

$$\begin{aligned} R_{\rm o} = k^2 \frac{1+q}{q}\Big(\frac{0.49 + 0.15q^{\frac{2}{3}}}{0.6 + q^{\frac{2}{3}}\ln(1+q^{-1/3})}\Big)^2 \left(1 + q\times\right. \\ \left.\Big(\frac{(0.49 + 0.27q^{\frac{1}{3}} - 0.12q^{\frac{2}{3}})[0.6 + q^{\frac{2}{3}}\ln(1+q^{-\frac{1}{3}})]}{(0.49q^{-\frac{2}{3}} + 0.15)[0.6q^{\frac{2}{3}} + \ln(1+q^{\frac{1}{3}})]}\Big)^2\right), \end{aligned} \tag{9}$$

setting equation $R_{\rm o} = 1/3$, and solving (numerically) for q, we obtain the minimum mass ratio, $q_{\rm min,out}$, for stability of W UMa systems in which two components have filled the outer Roche lobes with the different values of k^2 and the relation between $q_{\rm min,out}$ and k^2 is also shown in Figure 1. As seen from Figure 1, the value of $q_{\rm min,in}$ is smaller than that of $q_{\rm min,out}$ at any value of k^2. This suggests that the stability of a contact binary with given masses indeed depends on the degree of contact f, or filling factor, $0 < F = 1 - f < 1$. Meanwhile, the more is the value of k^2, the larger are the minimum mass ratios ($q_{\rm min,in}$ and $q_{\rm min,out}$) and the difference between $q_{\rm min,in}$ and $q_{\rm min,out}$.



Figure 1. The minimum mass ratio for stability $q_{\rm min}$ in the Roche approximation. The parameter k^2 is the dimensionless gyration radius of the stars

We take $k^2 = 0.06$ as Rasio (1995), the ratio of the spin angular momentum to the orbital angular momentum against the mass ratio is plotted in Figure 2. Meanwhile, we obtain $q_{\rm min,in} = 0.071$ and $q_{\rm min,out} = 0.078$ for $k^2 = 0.06$. In general, the dynamical instability of W UMa contact binary systems should be occurred at an overcontact stage with a higher degree of contact rather than at a marginal contact one. The numerical study of the equilibrium and stability of close binary systems (Rasio & Shaprio 1995) had identified the dynamical stability limit at a contact degree of about 70%, corresponding to a minimum mass ratio of about 0.076. This suggests that the W UMa systems with mass ratios in a region of 0.076–0.078 would suffer Darwin's instability and then coalesce into the fast-rotating single stars (including FK Com stars and blue stragglers), implying that the W UMa systems with mass ratios $q \leq 0.076$ can not be observed. We collect the absolute parameters of some of W UMa systems (listed in Table 1) reported recently, together with those compiled by Maceroni & van't Veer (1996), Gazeas et al. (2005), and by Yakut & Eggleton (2005), to determine the minimum mass ratio of the observed W UMa systems. Taking $k^2 = 0.06$



Figure 2. The angular momentum ratio R against the mass ratio, q. The solid lines represent theoretical results corresponding to the systems filling the inner Roche lobe and the outer Roche lobe, respectively. The solid dots represent the observed W UMa systems.



Figure 3. The angular momentum ratio R against the mass ratio, q. The solid line represents the fitting result. The solid dots represent the observed W UMa systems.

as mentioned above, the observed systems are also plotted in Figure 2. It is seen in Figure that, except for two systems with a larger scatter, all systems load in the region limited by the theoretical curves corresponding to the inner and outer Roche lobes, and none of them has surpassed the dynamical stability limit. However, it is difficult to obtain a direct polynomial relation between the ratio R and the mass ratio q by the least-squares fitting. But we have found a relation between the logarithms of the ratio R and the mass ratio q as the following

$$\log R = -1.4347 + 0.09954\log q + 1.8506(\log q)^2 + 1.25287(\log q)^3 + 0.3291(\log q)^4, \tag{10}$$

the observed systems, together with the theoretical curve, are shown in Figure 3. As seen from Figure 3, the theoretical curve has given a good fit to the observed systems. Setting equation $\log R = \log(1/3)$, and solving (numerically) for q, we obtain a minimum mass ratio, $q_{\rm min,obs} = 0.077$, for the observed systems. It is larger than 0.076 estimated from the theoretical result (Rasio & Shaprio 1995). This suggests that the observed W UMa systems have probably suffered the loss of angular momentum. Since the loss of angular momentum of the observed systems would result in the shrinking of their orbits and thus would cause their inner and outer Roche lobes to decrease and contact degree f to increase, so that leads f to have exceeded 70% before the observed systems reaching the minimum mass ratio $q = 0.076$.

Table 1. The physical parameters of some W UMa systems.

Stars	P (days)	M_1 ($M_\odot$)	q	R_1 ($R_\odot$)	R_2 ($R_\odot$)	Refs.
V410 Aur	0.3663	1.30	0.146	1.40	0.61	(1)
XY Boo	0.3705	0.91	0.186	1.23	0.61	(1)
AH Cnc	0.3605	1.21	0.149	1.36	0.62	(2)
CW Cas	0.3188	1.06	0.547	1.01	0.76	(3)
V776 Cas	0.4404	1.63	0.13	1.71	0.71	(4)
V899 Her	0.4212	2.10	0.566	1.57	1.22	(5)
BB Peg	0.3615	1.38	0.362	1.26	0.76	(6)
V351 Peg	0.5933	1.63	0.361	1.87	1.19	(7)
AU Ser	0.3865	0.90	0.710	1.10	0.94	(8)
VZ Psc	0.2613	0.81	0.800	0.78	0.70	(9)

References in Table 1:(1) Yang et al.(2005a); (2) Zhang et al.(2005); (3)Barone et al.(1988); (4) Djurasevic(2004); (5)Ozdemir et al. (2002); (6) Lu & Rucinski(1999); (7) Albayrak et al.(2005); (8) Gurol(2005); (9) Hrivnak et al.(1995)

3 DISCUSSIONS AND CONCLUSIONS

The models of low-mass W UMa-type systems (Li et al. 2004, 2005) have shown that W UMa systems would suffer Darwin's instability and evolve into fast-rotating single stars (i.e. FK Com stars or blue stragglers). Rasio (1995) predicted a cutoff of the mass ratio for W UMa systems at about 0.09. Considering the rotation of the secondary, we obtain that the minimum mass ratio for stability of the marginal contact binaries is of about $q_{\rm minin} = 0.071$, suggesting that the rotation of the secondary still has a definitely influence on the determination of the dynamical stability limit of W UMa systems. Rasio (1995) had predicted that the dynamical stability limit should depend on the fill-factor of W UMa systems. Using the new results in studies of the W UMa systems, we investigate the dynamical stability limit of W UMa systems in which both components fill their outer Roche lobes and obtain a minimum mass ratio $q_{\rm min,out}$ of about 0.078 for this kind of systems. This suggests that the dynamical stability limit of W UMa systems indeed depends on their fill-factor.

The difference between $q_{\rm min,in}$ and $q_{\rm min,out}$ suggests that the dynamical stability of the W UMa contact binaries with given mass depends on the degree of contact $[f = (\Omega - \Omega_{\rm in})/(\Omega_{\rm out} - \Omega_{\rm in})]$, which is consistent with the result predicted by Rasio (1995). Rasio & Shaprio (1995) had identified the dynamical stability limit of W UMa systems at about $f = 70\%$, which corresponds to a minimum mass ratio of about 0.076. We obtain a minimum mass ratio of about 0.077 , corresponding to a contact degree $f = (q_{\rm min,obs} - q_{\rm min,in})/(q_{\rm min,out} - q_{\rm min,in}) \approx 86\%$, for the observed systems. The difference between the Rasio & Shaprio's result and ours derived from the observed systems suggests that the observed W UMa systems have probably suffered the loss of angular momentum owing to GR or MSW during the evolution of the observed W UMa systems. In fact, *Einstein* soft X-ray observations and *International Ultraviolet Explorer (IUE)* Ultraviolet observations (Vaiana 1980; Eaton 1983; Rucinski & Vilhu 1983; Vilhu 1983) have shown that W UMa systems are strong sources. This suggests surface activity of the kind observed on our Sun, and so presence of the magnetic field. Since the observed W UMa systems rapidly lose the angular momentum via MSW, the contact degrees of them have exceeded 70% before their mass ratios decrease to the minimum mass ratio q=0.076.

Among the about 200 contact binaries with reliable photometric data, the system V857 Her has an extremely small mass ratio, $q = 0.0653$ (Qian et al. 2005) for observed W UMa systems at present. This is not only a new "record" for contact binary, surpassing the well-known AW UMa with $q = 0.075$ (Rucinski 1992) and SV Crv with $q = 0.066$ (Rucinski et al. 2001). It is a tremendous contribution for theory that W UMa systems with the extremely small mass ratios are found. Only a slight difference between observations and theory not only indicates that theory have come close to a decent prediction, but also indicates that some physical processes taking place in the observed W UMa systems have not be considered in the present theory. At first, the dynamical stability limit of W UMa systems may also depend on the structure (i.e. the value of k^2) and the value of k^2 may decrease with the evolution. In addition, the differential rotation of the observed systems would lead the ratio of the gyration radii to the stellar radii of the components of the W UMa systems to be decreased. Therefore, the differential rotation of the components of the W UMa systems is another physical reason why some W UMa systems with the mass ratios smaller than the minimum mass ratio 0.076 still can be observed. However, contact binaries with mass ratios significantly below that of SX Crv and V857 Her should not be observed, since they are tidally unstable and quickly merge into a single, fast-rotating object, on a tidal timescale of about 10^3–10^4 yr (Rasio 1995).

ACKNOWLEDGEMENTS

This work was partly supported by the Chinese Natural Science Foundations (Nos. 10273020, 10303006, and 10433030), the Yunnan Natural Science Foundation (Grant No. 2005A0035Q), and by the Foundation of Chinese Academy Sciences (KJCX2-SW-T06). We would like to thank the referee Prof. F. A. Rasio for his helpful comments and suggestions which greatly improved this paper.

REFERENCES

Albayrak B., Djurasevic G., Selam S. O., et al., 2005 New Astron., 10, 163
Barone F., Milano L., Maceroni C., & Russo G., 1988, A&A, 197, 347
Djurasevic G., Albayrak B., Selam S. O., et al., 2004, New Astron., 9, 425
Eaton J. A., 1983, ApJ, 268, 800
Eggleton P. P. 1983, ApJ, 268, 368
Eggleton P. P., & Kiseleva-Eggleton L., 2001, ApJ, 562, 1012
Gazeas K. D., Baran A., Niarchos P., Zola S., et al. 2005, Acta Astron., 55, 123
Gurol B., 2005, New Astron., 10, 653
Hut P., 1980, A&A, 92, 167
Hrivnak B. J., Guinan E. F., & Lu W. X., 1995, ApJ, 455, 300
Kaluzny J. 1990, Acta Astron. 40, 61
Kaluzny J., & Shara M. M., 1988, AJ, 95, 785
Li L., Han Z., & Zhang F., 2004, MNRAS, 355, 1383
Li L., Han Z., & Zhang F., 2005, MNRAS, 360, 272
Lu W. X., & Rucinski S. M., 1999, AJ, 118, 515
Maceroni C., & van't Veer F., 1996, A&A, 311, 523
Mateo M. 1996, in *The Origins, Evolutions, and Destinies of Binary Stars in Clusters*, ASP Conf., Vol. 90, ed. E. F. Milone & J. C. Mermilliod, 346
Mateo M., Harris H. C., Nemec J., & Olszewski E. W., 1990, AJ, 100, 469
Ozdemir S., Demircan O., Erdem A. et al., 2002, A&A, 387, 240
Qian S., Zhu L., Soonthornthum B., Yuan J., et al., 2005, AJ, 130, 1206
Rasio F. A. 1995, ApJ, 444, L41
Rasio F. A. & Shapiro S. L., 1995, ApJ, 438, 887
Rucinski S. M., 1992, AJ, 104, 1968
Rucinski S. M., Lu W. X., Mochanacki S. W., Ogloza W., & Stachowski G., 2001, AJ, 122, 1974
Rucinski S. W., Vilhu O., 1983, MNRAS, 202, 1221
Vaiana G., 1980, in Wayman P.A., ed., Highlights of Astronomy, Vol. 5, Reidel, Dordrecht, P. 419
Vilhu O., 1983, in West R. M., ed., Highlights of Astronomy, Vol. 6. Reidel, Dordrecht, P. 643
Yakut K., & Eggleton P. P., 2005, ApJ, 629, 1055
Yan L., & Mateo M., 1994, AJ, 108, 1810
Yang Y.-G., Qian S.-B., & Zhu L.-Y., 2005a, AJ, 130, 2252
Yang Y.-G., 2005b, PhD Thesis, Yunnan Observatory of Chinese Academy Sciences.
Zhang X. B., Zhang R. X., & Deng L., 2005, AJ, 129, 979

This paper has been typeset from a TEX/ LATEX file prepared by the author.